About Autism



CHILDREN ARE DIAGNOSED ON THE AUTISM SPECTRUM

Pediatrics: December 2018, VOLUME 142 / ISSUE 6: The Prevalence of Parent-Reported Autism Spectrum Disorder Among US Children.

Risk Factors identified by the Centers of Disease Control and Prevention (2019):

- Most scientists agree that genes are one of the risk factors that can make a person more likely to develop ASD.
- Children who have a sibling with ASD are at a higher risk of also having ASD.
- ASD tends to occur more often in people who have certain genetic or chromosomal conditions, such as fragile X syndrome or tuberous sclerosis External.
- When taken during pregnancy, the prescription drugs valproic acid and thalidomide have been linked with a higher risk of ASD.
- There is some evidence that the critical period for developing ASD occurs before, during, and immediately after birth.
- Children born to older parents are at greater risk for having ASD.

https://www.cdc.gov/ncbddd/autism/research.html.



43.3%

received behavior-based treatment

Most children with current ASD:

6.9%

received medication treatment only

JAMA Pediatrics | Original Investigation (2019). "Prevalence and Treatment Patterns of Autism Spectrum Disorder in the United States, 2016".



20.3%

received both behavioral and medication

Organizations that endorse behavioral treatment for children diagnosed with ASD:

- United States Surgeon General
- American Academy of Neurology
- American Academy of Family Pediatrics
- American Academy of Pediatrics
- American Psychological Association
- Society for Developmental and Behavioral Pediatrics
- Autism Society of America
- National Institute of Child Health & Human Development
- · National Institute of Mental Health



Numerous other outcome studies show substantial gains in:

- Measured IQ
- Educational Achievement
- Adaptive Behavior

(Lovass, 1986)



ButterflyEffects

Transforming Lives Through ABA Therapy

www.butterflyeffects.com (888) 880-9270

